

# Prevalence and Associated Factors of Diaper Rash Among Infants and Toddlers Aged 2 to 24 Months in Ola During Children's Hospital, University of Sierra Leone Teaching Hospital Complex, Freetown, Sierra Leone

<sup>1</sup>Ofovwe Gabriel E., <sup>1</sup>Bell Nellie T., <sup>2</sup>Ikhurionan Paul E.

<sup>1</sup>Ola During Children's Hospital, Sierra Leone Teaching Hospital Complex, Freetown, Sierra Leone.

<sup>2</sup>Department of Child Health, University of Benin Teaching Hospital, Benin-City, Edo State Nigeria.

Corresponding Author: Prof Ofovwe G.E.; ofowwegabriel@yahoo.com

**How to Cite:** Ofovwe G. E., Bell N. T. & Ikhurionan P. E. Prevalence and associated factors of diaper rash among infants and toddlers in Ola During Children's Hospital, University of Sierra Leone Teaching Hospital Complex, Freetown, Sierra Leone: SLJM 2024;Vol 1(1) pp 1-4

**Editor:** Prof. Kehinde S. Oluwadiya, University of Sierra Leone Teaching Hospitals Complex, Freetown, Sierra Leone

**Received:** May 2023

**Accepted:** November 2023

**Published:** March 2024

**Copyright:** © 2024, Ofovwe Gabriel. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

**Funding:** No funding was received for this study.

**Conflict of interest:** The authors declare they have no conflicts of interest that are directly or indirectly related to the research.

## ABSTRACT

**Background:** Diaper rash is a common dermatological condition in infant and toddlers. This condition is also commonly seen particularly among children with malnutrition at Ola During Children's Hospital (ODCH), Freetown, Sierra Leone. However, the prevalence of this condition in Sierra Leone is not known. This study therefore was conducted to determine the prevalence of prevalence diaper rash and associated factors in infants and toddlers admitted to ODCH.

**Methods:** This was a descriptive cross-sectional study of 269 children aged 2 to 24 months. A convenience sampling technique was used to select the subjects after verbal informed consent was obtained. A structured questionnaire was used to collect data. All children were examined, and a description of the perineum was also documented at the time of recruitment. Analysis was done with IBM SPSS version 26.0.

**Results:** The prevalence of diaper rash was 52%. The proportion of males with diaper rash compared to females was not statistically different. Artificial milk, cereal, malnutrition and immediate caregivers other than fathers were the factors significantly associated with diaper rash.

**Conclusion:** The prevalence of diaper rash is high among infants and toddlers seen at the ODCH, Freetown, Sierra Leone. Factors associated with diaper rash include type of diet, nutritional status and immediate caregiver.

**Key Words:** Prevalence, associated factors, diaper rash, Freetown.

## INTRODUCTION

Diaper rash is a common skin condition in children below 24 months globally.<sup>1</sup> It results from prolong contact between the area of the skin covered by diaper and the diaper soiled with urine and faeces.<sup>2</sup> The production of ammonia from urinary urea by faecal bacteria increases skin pH and thus promotes skin susceptibility to mechanical damage and the activity of faecal enzymes such as proteases and lipases.<sup>2</sup> These acting together result in irritant contact dermatitis often characterized by erythematous and scaly skin, papulovesicular or bullous lesions, fissures, and erosions. The prevalence of diaper rash varies from country to country and from region to region even within the same country.<sup>3</sup> In Italy, 15% was reported whereas in Japan the prevalence is as high as 87%.<sup>3</sup> Similarly, studies from Africa have reported varying prevalence ranging from 2.7% in Ibadan, Nigeria<sup>4</sup> to 18.4% in Yaoundé, Cameroon<sup>5</sup> and 62.5% in Adama, Ethiopia.<sup>6</sup>

Certain factors have been reported to be associated with or contribute to the development of diaper rash. These factors vary according to setting, hygiene practices, and age group of the child. Commonly identified factors include the non-use of disposable diapers, low socioeconomic status, lack of breastfeeding, diaper care practices such as infrequent changes and use of barrier creams.<sup>7,8</sup> Environmental factors such as humidity and temperature have also been suggested to play a role in the development of diaper rash.<sup>3</sup> Whereas, diaper rash was observed to be common in infants particularly malnourished children admitted to Ola During Children's Hospital, Freetown, very little is known about the prevalence and factors associated with the condition in Sierra Leone. These children might have specific risk factors for diaper dermatitis due to differences in skin sensitivities. This study was therefore undertaken to determine the prevalence and factors associated with the condition in this locale.

## METHODS

This was a descriptive cross-sectional study conducted over a 14-week period, 1st December 2021 to 15th March 2022 at Ola During Children's Hospital (ODCH), University of Sierra Leone Teaching Hospital Complex, Fourah Bay Road, Freetown, Sierra Leone. ODCH is located in the densely populated eastern part of Freetown, Sierra Leone. It is the only tertiary hospital for sick children and is a part of the University of Sierra Leone Teaching Hospital Complex, which comprises of six hospitals. It receives referrals from across the country. It has a bed capacity of 197. It comprises of an emergency ward (for resuscitation and stabilization of patients), high dependency ward, intensive care unit (ICU) 3 main wards, special care baby unit, general and specialist outpatient clinics. Admission into the hospital is usually through the emergency unit. The general out-patient clinic attends to children whose conditions do not meet admission crite-

ria (such as common cold, uncomplicated malaria, diarrhea diseases with no dehydration etc.) and hence managed on out-patient bases. The general out-patient clinic is staffed by resident doctors and house officers who rotate through the emergency unit. These doctors are supervised by a consultant. An average of 20 to 40 children is attended to in the out-patient clinic daily.

### Subjects and procedure:

The subjects comprised of children aged two months to twenty-four months, who were seen as outpatients in the emergency ward and those admitted for various conditions in the hospital. A convenience sampling technique was used to select the subjects. A 32-item questionnaire was drawn up and administered to parents of children wearing diaper who were either admitted in the hospital or seen in the outpatient clinic. The questionnaire consists of different variables such as socioeconomic, demographic data and factors associated with diaper rash. The questionnaire was structured with simple to complete questions. Diaper dermatitis was defined as a rash in the diaper region thought to be caused by the child wearing diapers. Demographic data (age, location, primary care giver and parents' marital status); history of allergic conditions (and family history), number of diarrhea episodes in the preceding six months, diet and nappy use were documented. Frequency of diaper change was considered good if it was at least eight times a day and poor if less than eight times a day. Nutrition status was considered to be well-nourished if the weight-for-age was greater than -2 Z-scores while weight-for age less than -2 Z-score was judged as malnourished. All children had a full examination including anthropometric measurements (weight and length) as well as description of the perineum at the time of recruitment. The questionnaires were completed by the researchers and trained research assistants.

### Statistical analysis:

The collected data were organized, tabulated, and statistically analyzed using the International Business Machines Corporation (IBM) SPSS version 26.0 (SPSS for Window Inc; Chicago, LL, USA) Statistical Software. Descriptive statistics such as frequency, percentage, mean and standard deviation were used to describe independent variables. Chi-square test was used to establish the association between dependent and independent variables. Multivariate regression analysis was used to identify factors that were independently associated with diaper rash. A p-value of less than 0.05 was accepted as the level of statistical significance.

### Ethics and consent:

Informed verbal consent to participate in the study was obtained from primary care giver of all subjects after explanation of the study objectives, relevance, risks, and benefits. Permission for the study was obtained from the hospital's authority. Participation in the study was entirely voluntary and participants were advised that they could withdraw from the study at any point.

## RESULTS

### Socio-demographic characteristics:

A total of two hundred and sixty-nine subjects were recruited comprising of 155 (57.6%) males and 114 (42.4%) females. The mean  $\pm$  SD age of the subjects was  $8.1 \pm 0.4$  months. Majority, 208 (77.3%) of the subjects were infants while 61 (22.7%) were aged 13 – 24 months. Majority of the mothers 165 (61.3%) were not married while 68.4% were aged between 20 and 30 years. The distribution of other sociodemographic factors such as educational level, immediate caregivers and level of income of the family are as shown in Table I.

### Prevalence of Diaper rash:

More than half of the study population (140/269, 52.8%) had diaper rash at the time of data; Sixty-eight (25.3%) had erythema in diaper area, 46 (17.1%) with hypo-pigmented lesions, 14 (5.2%) with papules/pustules, while 12 (4.5%) had excoria-

tions/ulcerated lesions.

**Table I:** Distribution of sociodemographic characteristics of the mothers and subjects

Variable	n	%
<b>Child's age group</b>		
Infants (2 – 12 months)	208	77.3
Older child (13 – 24 months)	61	22.7
<b>Sex</b>		
Male	155	57.6
Female	114	42.4
<b>Mothers' age group</b>		
Teenagers ( $\leq 19$ yrs)	29	10.8
20 – 30 yrs	184	68.4
$\geq 30$ yrs	56	20.8
<b>Immediate caregiver</b>		
Mother	170	63.2
Father	33	12.3
Grandmother	17	6.3
Both parents	43	16.0
Others (sibling, aunty)	6	2.2
<b>Mothers' educational level</b>		
Primary/none	145	53.9
Secondary	106	39.4
Tertiary	18	6.7
<b>Mothers' marital status</b>		
Married	104	38.7
Not Married	165	61.3
<b>Family monthly income</b>		
< \$50	215	79.9
\$50 - \$100	46	17.1
> \$ 100	8	3.0

### Factors associated with diaper rash:

About equal proportion of males 81/155 (52.3%) compared to females 59/114 (51.8%) had diaper rash,  $\chi^2 = 0.007$ ;  $P > 0.05$ . Similarly, there was no difference between the proportion of infants with diaper rash 107/208 51.4% and those aged between 13 and 24 months 33/61, 54.1%,  $\chi^2 = 0.133$ ;  $P > 0.05$ . Table 2 shows the relationship between the prevalence of diaper rash and other factors.

### Multivariable Regression of factors associated with diaper rash:

In a multivariate regression analysis showed that only nutritional status was independently associated with diaper rash in children (Table III).

## DISCUSSION

This study sought to determine the prevalence and associated factors of diaper rash in children aged two months to 24 months. We found diaper rash in over half (52%) of the children recruited. The presence of diaper rash was associated with childhood malnutrition, consumption of formula feeds and cereal-based food. Surprisingly, children whose primary caregivers were the fathers were less likely to have diaper rash compared to those cared for by mothers or other relatives. The prevalence of diaper rash seen in the present study was higher than that reported in Thailand (36.1%),<sup>2</sup> Saudi Arabia (39.3%),<sup>9</sup> Port Harcourt, Nigeria (34%)<sup>10</sup> and Yaoundé, Cameroon (18.4%).<sup>5</sup> Our finding is however lower than the prevalence described in Ethiopia (62.5%)<sup>6</sup> and Mauritius (79.7%).<sup>3</sup> The difference in prevalence observed in our study may be due to climatic difference and hygiene practices in the study population.

Nutritional status was associated with diaper rash as the proportion of children with malnutrition who had diaper rash was significantly higher than the proportion in well-nourished children. The association between nutritional status and diaper rash may be due to the fact that malnourished children are commonly deficient in some nutrients including zinc which plays an important role in maintaining the integrity of the skin and immunity. Deficiency of zinc may thus predispose to diaper dermatitis in the presence of minimum friction and irritation.

Furthermore, the impaired immunity associated with severe malnutrition predisposes to infections with bacteria such as

**Table II:** Age specific prevalence and factors associated with diaper rash among infants and toddlers in ODCH

Factors	Diaper rash		$\chi^2$ - value	P - value
	Present n (%)	Absent n (%)		
<b>Child's age group (months)</b>				
2 – 6	57 (47.9)	62 (52.1)	2.119	0.548
7 – 12	49 (56.9)	37 (43.1)		
13 – 18	22 (56.4)	17 (43.6)		
> 18	12 (48.0)	13 (52.0)		
<b>Sex</b>				
Male	81 (52.3)	74 (47.7)	0.007	0.935
Female	59 (51.8)	55 (48.2)		
<b>Mother's age group</b>				
≤19 yrs.	18 (60.1)	11 (39.9)	2.430	0.297
20 – 30 yrs.	97 (52.7)	87 (47.3)		
≥30 yrs.	25 (44.6)	31 (55.4)		
<b>Mother's educational level</b>				
None/Primary	75 (51.7)	70 (48.3)	0.096	0.953
Secondary	55 (51.9)	51 (48.1)		
Tertiary	10 (55.6)	8 (44.4)		
<b>Immediate caregiver</b>				
Mother	112 (54.9)	92 (45.1)	7.144	0.028*
Father	10 (30.3)	23 (69.7)		
Others <sup>+</sup>	18 (56.3)	14 (43.7)		
<b>Child's diet</b>				
Breast milk only	14 (33.3)	28 (66.7)	9.889	0.04*
Breast & artificial milk	48 (53.3)	42 (46.7)		
Cereal	40 (64.5)	22 (35.5)		
Mixed diet	17 (51.5)	16 (48.5)		
Family diet	21 (50.0)	21 (50.0)		
<b>Frequent diaper changes</b>				
Good	8 (40.0)	12 (60.0)	1.256	0.262
Poor	132 (53.0)	117 (47.0)		
<b>Type of diaper</b>				
Disposable only	75 (46.9)	85 (53.1)	4.229	0.121
Re-usable towel only	12 (60.0)	8 (40.0)		
Both	53 (59.6)	36 (40.4)		
<b>Child's nutritional status</b>				
Well nourished	66 (45.2)	80 (54.8)	5.984	0.01*
Malnourished	74 (60.2)	49 (39.8)		
<b>Number of diarrhoea episodes</b>				
0	103 (50.7)	100 (49.3)	0.659	0.719
1	18 (58.1)	13 (41.9)		
≥ 2	19 (54.3)	16 (45.7)		

\* - Statistically significant; + - Grandmother, aunty, siblings

Staphylococcus aureus and fungi such as Candida albicans, two organisms reported to be associated with diaper rash.<sup>7</sup> It is not surprising therefore that diaper rash was more prevalent in children with malnutrition. Previous studies had not considered the impact of nutritional status on the prevalence of diaper rash. Also, children who are malnourished are more likely to be from the low socioeconomic strata and thus may not afford disposable diapers or change them frequently.

Type of diet also plays a significant role in development of diaper rash as cereal-based diet was significantly associated with diaper rash. The consumption of artificial milk in addition to breast milk or as mixed diet also contributes to development of diaper rash as more children exposed to these types of diet

when transitioning babies from exclusive breast milk to cereal-based meals.

Immediate caregiver is also significantly associated with development of diaper rash as care given by mothers, grandmothers, aunts, siblings and others is significantly associated with development of diaper rash. Surprisingly, care given by fathers is less likely to result in diaper rash. It is likely that children who were cared for primarily by their fathers were less likely to use re-usable towels than those catered for by mothers and other relatives. The association between use of cloth diapers and development of diaper rash has been shown by previous studies. Although our study did not show significant association between diaper rash and type of diaper used, subjects who were diapered with re-usable towel alone or in com-

**Table III:** Multivariable Regression of factors associated with diaper rash

Variable	B	Wald	OR	95% CI	p
Child's diet	-0.064	7.791	0.938	0.773-1.138	0.517
Immediate care giver	-0.476	3.347	0.067	0.373-1.035	0.067
Nutritional status	-0.525	4.216	0.591	0.358-0.976	0.040*

\*significant at  $p < 0.05$ ; OR – odds ratio; CI – confidence interval;

developed diaper rash than those who did not. Previous studies have reported the effect of diet on diaper rash and suggested that babies who were breast-fed exclusively have stools of lower pH and fewer urease producing bacteria and lower incidence of diaper rash.<sup>3,11</sup> The practice of exclusive breastfeeding in Freetown has previously been reported to be poor particularly among young and unmarried mothers.<sup>12</sup> Majority of mothers in this study were unmarried. The increase in stool pH with introduction of cereals (either alone or in combination with breast milk) increases the predisposition to nappy rash. It is therefore necessary to pay more attention to diaper hygiene and change which prevent diaper rash especially

when transitioning babies from exclusive breast milk to cereal-based meals. bination with disposable diapers had the higher proportion of diaper rash compared to disposable alone (60% and 59.6% respectively compared to 46.9). It is possible that since re-usable towels are not disposable; they are not frequently changed thereby staying longer periods on the children than disposable diapers. The study also showed that teenage mothers and less frequent diaper changes (less than 8 times per day) were predisposing factors for diaper rash, although the association wasn't statistically significant. Poor diaper practice has been previously reported to be associated with diaper rash.<sup>11</sup> The reasons are not farfetched as infrequent diaper changes will lead to prolonged contact of soiled diaper with the

skin and increase irritation whereas, teenage mothers are more likely to lack the skills required for care of their babies. It is to be noted that teenage pregnancy and unmarried mothers are common in Sierra Leone.

## CONCLUSION

Prevalence of diaper rash among children two months to 24 months in Freetown is high. While malnutrition is significantly associated with its occurrence, other predisposing factors include diet type particularly cereal and artificial milk and immediate care giver are shown to also be associated. Poverty and lack of knowledge appear to be underlying causes of these factors. It is recommended that basic health education on personal hygiene and diaper care as well as strengthening of breastfeeding practices should be provided for mothers during antenatal and immunization visits.

## LIMITATION OF THE STUDY

Since this is a hospital-based study, it may not adequately represent what is happening in the community. Furthermore, the study may also be affected by social-desirability bias being a questionnaire based study. Also, the use of convenient (non-probability) method for sampling might have inadvertently excluded or over-represented children with diaper dermatitis.

## ACKNOWLEDGEMENT

We acknowledge and thank the mothers who participated in the study and the research assistants for their help in administering the questionnaire.

## CONFLICT OF INTEREST

All authors declare that we do not have any financial or personal relationship which may have inappropriately influenced us in writing this article.

## REFERENCE

- Collins G, Inusah AW, Asumah MN, Kwarteng PG, Ziblim SD, Dzomeku P. Knowledge of mothers with children age 1 to 24 months on diaper dermatitis management and associated practice in a referral hospital in Northern Ghana: a cross sectional study. *Pan African Med J One Heal*. 2022; 8:1-15.
- Sukhneewat C, Chaiyarit J, Techasatian L. Diaper dermatitis: A survey of risk factors in Thai children aged under 24 months. *BMC Dermatol*. 2019; 19(1):1752-1760.
- Biranjia-hurdoyal S, Pandamikum L. A Study to Investigate the Prevalence of Nappy Rash among Babies Aged 0 to 36 Months Old in a Tropical Country. *Austin J Dermatolog*. 2015; 2(2):1040.
- Owa AB, Oladokun RE, Osinusi K. Diaper dermatitis among children in Ibadan, Nigeria: frequency and predisposing factors. *Eur J Pediatr Dermatol*. 2016; 26(3):135-141.
- Chiabi A, Ngamgo Kamdem JC, Nkoro AG, Siyou H, Mah E, Nguetack FD et al. Epidemiological and Clinical Features of Diaper Dermatitis in Infants at the Yaounde Gynaeco-Obstetric and Pediatric Hospital, Cameroon. *J Child Sci*. 2018; 8(1):e46-e49.
- Jewaro S, Mariam YG, Ali B. Prevalence of diaper rash and associated factors among babies aged 0 to 24 months in Adama Administrative City, Central Ethiopia: A cross sectional study. *Afr J Med Heal Sci*. 2021;20(5):53-61.
- Jordan WE, Lawson KD, Berg RW, Franxman JJ, Marner AM. Diaper Dermatitis: Frequency and Severity Among a General Infant Population. *Pediatr Dermatol*. 1986; 3(3):198-207.
- Roy S, Mollah MAH, Mannan MA, Khan ZJ, Al Amin A, Ghosal S et al. Hazards of Wearing Diaper among Infants. *J Bangladesh Coll Physicians Surg*. 2017; 35(4):170-173.
- Alghamdi MA, AL-Ghamdi HS, AlHajji AM, Alzahrani SA, Al-Thobaiti LY, Alzahrani SA et al. Prevalence and risk factors of diaper dermatitis among newborn babies to two years of age in Al-Baha region, Saudi Arabia. *Middle East J Fam Med*. 2022; 7(10):47-57.
- Eke GK, Opara P Ibo. Mothers Knowledge and Home Management of Nappy Rash in Port Harcourt, Nigeria. *Niger Heal J*. 2013; 13(4):152-157.
- Siraj Jewaro, Yemane G. Mariam and Belay Ali. Prevalence of diaper rash and associated factors among babies aged 0 to 24 months in Adama Administrative City, Central Ethiopia: A cross sectional study. *Afr. J. Med. Health Sci*. 2021; 20 (5): 53-61.
- Bell NVT, Ikhurionan PE, Akhigbe IE, Mustapha AM and Ofofwe GE. Knowledge attitude and practices of breastfeeding among mothers of children six months to three years seen at a tertiary children hospital in Freetown. *Niger J Paediatr* 2022; 49(2): 109 – 116.